Table 8. Estimated numbers of persons living with HIV/AIDS by year and selected characteristics, 1999–2002—30 areas with confidential name-based HIV infection reporting

	1999	2000	2001	2002	
Age as of end of year (yrs)					
<13	2,517	2,488	2,451	2,363	
13—14	242	291	394	467	
15—24	8,984	9,440	9,901	10,295	
25—34	59,153	57,347	56,314	55,789	
35—44	98,193	106,376	112,845	118,714	
45—54	45,171	53,002	61,547	70,472	
55—64	10,732	12,830	15,240	18,591	
≥65	2,985	3,635	4,407	5,241	
Race/ethnicity					
White, not Hispanic	91,038	96,424	101,943	107,992	
Black, not Hispanic	112,483	121,903	131,360	141,184	
Hispanic	21,256	23,449	25,732	28,364	
Asian/Pacific Islander	791	923	1,045	1,181	
American Indian/Alaska Native	1,224	1,339	1,431	1,565	
Exposure category					
Male adult or adolescent					
Male-to-male sexual contact	100,582	108,174	116,250	125,268	
Injection drug use	31,554	32,880	34,093	35,380	
Male-to-male sexual contact and injection drug use	14,900	15,290	15,696	16,143	
Heterosexual contact	19,594	22,021	24,387	26,843	
Othera	2,673	2,735	2,819	2,922	
Subtotal	169,304	181,099	193,244	206,557	
Female adult or adolescent					
Injection drug use	16,666	17,504	18,209	18,831	
Heterosexual contact	37,657	42,239	46,841	51,538	
Other ^a	1,306	1,414	1,513	1,627	
Subtotal	55,629	61,158	66,563	71,996	
Child(<13 yrs)					
Perinatal	2,741	2,855	3,006	3,114	
Other ^b	297	291	279	260	
Subtotal	3,038	3,146	3,285	3,374	
Total ^c	227,976	245,409	263,098	281,931	

Note. These numbers do not represent actual cases in persons living with HIV/AIDS. Rather, these numbers are point estimates of persons living with HIV/AIDS that have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk. The estimates have not been adjusted for incomplete reporting.

Data include persons with a diagnosis of HIV infection. This includes persons with a diagnosis of HIV infection only, a diagnosis of HIV infection and a later AIDS diagnosis, and concurrent diagnoses of HIV infection and AIDS.

Since 1998, the following 30 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Utah, Virginia, West Virginia, Wisconsin, Wyoming, and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting only for new diagnoses.

^a Includes hemophilia, blood transfusion, perinatal, and risk not reported or not identified.

^b Includes hemophilia, blood transfusion, and risk not reported or not identified.

^c Includes persons of unknown or multiple race and of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.